

Index

- affine
 - dimension, 29
 - hull, 29
 - independence, 29
- algorithmic parameters (tuning of), 3
- analysis of variance, 6, 234
- automatic differentiation, 2
- automatic error analysis, 3

- backtracking (line search), 163, 164, 168
- barrier
 - extreme barrier function, 242, 243, 245, 247
- benchmarking, *see* data profiles
- black box, 2, 242, 249

- Cauchy step, *see* step
- circuit design, 4
- Clarke
 - generalized directional derivative, 125
 - nonsmooth analysis, 125, 127, 247
 - strict differentiability, 127
- combinatorial optimization, 6
- completing
 - a nonpoised set
 - via Lagrange polynomials, 95, 101–102
 - via LU factorization, 99–102
- condition number, 4, 99, 225, 231
 - as a measure of well posedness, 38–39, 48–52, 68–69, 84–86
 - in error bounds, 52, 54, 69, 79

- cone
 - normal, 244
 - of descent directions, 125
 - polar, 244
 - positively spanned by vectors, 15–19, 244
 - tangent, 244, 246
- constraints
 - ϵ -active, 244
 - active, 244, 245
 - hidden, 242, 247
 - linear, 241, 244–248
 - relaxable, 241–242, 246–248
 - simple bounds, 241, 243–244, 247
 - unrelaxable, 241–242, 247, 248
 - with derivatives, 243–246, 248
 - without derivatives, 246–249
- convergence
 - global (definition), 120
 - global (results), 122, 124, 126, 131, 133, 155, 157, 167, 189, 211, 214, 219
 - global (results for second-order critical points), 198, 211, 225
 - lack of (Nelder–Mead), 8, 10, 149
 - local, xi, 5, 35, 74, 182
- convex, *see also* cone
 - hull, 30, 41
- coordinate search, *see* methods
- cosine measure, *see* positive spanning set
- Cramer’s rule, 41, 71, 77, 82
- criticality step, 165, 166, 183–185, 192–193, 195

- curvature, 9–11, 35, 73, 74, 80, 135,
136, 141, 145, 173, 174, 231
negative, 177, 178, 191
positive, 175
- data profiles, 9
- decomposition, *see* factorization
- decrease
imposing sufficient decrease in
directional direct search,
134–135
imposing sufficient decrease in line
search, 164
imposing sufficient decrease in
modified Nelder–Mead,
150–153
lexicographic (Nelder–Mead),
144–145
simple, 116, 119, 131, 132, 139,
150, 162, 178, 182, 183, 191,
204, 208, 215, 236, 245
sufficient, 119, 121, 135, 136, 139,
159–161, 163, 165–169, 176,
178, 182, 208, 211, 218, 236,
237, 245–247, *see also*
fraction of Cauchy decrease;
fraction of eigenstep decrease;
fraction of optimal decrease
- derivatives, *see also* automatic
differentiation; simplex
derivatives; simplex gradient;
simplex Hessian
finite differences, 2, 230
generalized finite differences, 47
- descent direction, 15
- dynamic pricing, 5
- eigenstep, *see* step
- engineering design, 4
- error bounds
for centered simplex gradients, 170
for polynomial interpolation
general case, 56
linear case, 26–28
quadratic case, 52–56
for polynomial regression
linear case, 28–29
quadratic case, 69–70
for simplex gradients, 33
for underdetermined quadratic
polynomial interpolation
(general), 78–80
for underdetermined quadratic
polynomial interpolation (in
the minimum Frobenius norm
sense), 83–84
in larger concentric balls, 200–202
Taylor-like, 35, 64, 69–70, 78, 83,
179
- factorization
by rows, 26
Cholesky, 20
eigenvalue decomposition, 203
Gaussian elimination, *see* LU
factorization
growth factor, 103, 104, 112
LDU, 112
LU, 99, 101–103, 105, 108
minimization of growth factor, 3–4
QR, 76, 245
singular value decomposition
(SVD), 29, 68, 76
- finite differences, *see* derivatives
- forcing function, 134, 151, 156, 159,
160, 236
- fraction of Cauchy decrease, 176, 220,
237–239
- fraction of eigenstep decrease, 177, 223,
238
- fraction of optimal decrease, 178, 205
- fully linear (FL), *see also* model
certificate/certifiable, 91, 180
class, 90, 180
- fully quadratic (FQ), *see also* model
certificate/certifiable, 91, 181
class, 91, 181
- function evaluations, 5, 10, *see also*
noise
cheap or inexpensive, 57, 229, 235

- costly or expensive, xi, 2, 4, 5, 8, 58, 70, 73, 92, 107, 178, 182, 183, 209, 229, 230, 242, 249
- noisy, 2, 8, 57, 70, 92
- generalized pattern search (GPS), *see* methods
- generating set search, 136–137
- geometry, *see also* completing; improvement; improving; poisedness
 - dispensing control of, 225
- global optimization, 249–250
- gradient, *see* derivatives; simplex
- groundwater community problems, 5
- group partial separability, 74
- Hessian, *see* derivatives; simplex
- heuristic methods
 - artificial neural networks, 6, 231
 - enumeration techniques, 6
 - evolutionary algorithms, 6
 - genetic algorithms, 6
 - particle swarm, 6, 249
 - population-based algorithms, 6
 - simulated annealing, 6
 - tabu-search, 6
 - variable neighborhood search, 250
- implicit filtering, *see* methods
- improvement, *see* model-improvement
 - algorithm; model-improving iteration; model-improvement step
- improving
 - well poisedness
 - minimum Frobenius norm models, 110
 - regression models, 108–110
 - via Lagrange polynomials, 95–98
 - via Lagrange polynomials (example), 98–99
 - via LU factorization, 104–106
 - via LU factorization (example), 106
- integer lattices, 123, 139, 151, 236, 245
 - globalization by, 127–130, 132, 135, 136, 139, 162, 245
 - tightness of integrality and rationality requirements, 130
- interpolation, *see also* error bounds; Lagrange polynomials; model; Newton fundamental polynomials; poisedness; uniqueness
 - conditions, 24, 37, 73
- Kriging, *see* model
- Lagrange polynomials
 - interpolation
 - basis for, 39
 - computing and updating, 40, 93–94
 - definition, 39
 - equivalent definition, 41
 - uniqueness, 39
 - volume definition, 41
 - invariant under scaling, 46, 212
 - invariant under shift, 46, 212
 - minimum Frobenius norm
 - computing and updating, 110
 - definition, 81
 - equivalent definition, 81
 - uniqueness, 87
 - volume definition, 82
 - minimum norm
 - basis for, 77
 - definition, 76
 - equivalent definition, 77
 - uniqueness, 77, 87
 - volume definition, 78
 - regression
 - basis for, 60
 - computing and updating, 108–109
 - definition, 60
 - equivalent definition, 62
 - uniqueness, 60
- lattice, *see* integer lattices
- Lebesgue constant, 40

- Lipschitz continuity
- of a function near a point
 - as an assumption, 126
 - of gradient
 - as an assumption, 23, 26, 28, 79, 90, 122, 166, 167, 179
 - of Hessian
 - as an assumption, 52, 69, 91, 170, 180
 - of model gradient, 92, 112
 - as an assumption, 90, 180
 - of model Hessian
 - as an assumption, 91, 181
- Lipschitzian optimization, 250
- manifold
- Lipschitz, 246
 - polynomial, 38
 - Riemannian, 246
- medical image, 5
- mesh adaptive direct search (MADS),
see methods
- mesh or grid
- assumptions (integer lattices), 128
 - coordinate search, 117
 - general, 127
 - minimal space between points, 128
 - other definitions, 130–131, 139
 - poll and search steps, 128, 132
 - search step, 118, 132
- methods
- augmented Lagrangian, 246–247
 - coordinate search, 9–11, 115–118, 125, 127, 135
 - directional direct search, 115–139, 235–237, 242–248
 - Dlviding RECTangles (DIRECT), 250
 - filter, 247–249
 - generalized pattern search, 118, 119
 - implicit filtering, 7, 10–11, 163, 168–170
 - interpolation-based trust region, 10–11, 204–205, 207–225, 248–249
 - last resort, 6
 - MADS (mesh adaptive direct search), 132–134, 136, 139, 247–248, 250
 - modified Nelder–Mead, 149–162
 - multidirectional search (MDS), 137–139, 150, 161, 162
 - multilevel coordinate search (MCS), 250
 - Nelder–Mead, 1, 7, 8, 10–11, 141–149, 161
 - parallel, 136, 168, 225, 248, 250
 - pattern search, 118, 123, 135, 136
 - quasi-Newton, 5, 10, 87, 168, 169, 204
 - sequential quadratic programming (SQP), 248–249
 - simplicial direct search, 137–139, 141–162, 237
 - trust region, 173, 225, 237–240, 248–249
 - using line search and simplex derivatives, 163–167
 - wedge (trust region), 215–225
- minimax optimization, 250
- mixed integer programming, 250
- model
- by space mapping, 239–240
 - fully linear (FL), 90, 180, 200–201, 210–214, 218–220
 - fully quadratic (FQ), 91, 181, 201–202, 210–211, 218–220
 - least Frobenius norm updating of quadratic models, 86, 212
 - model-improvement algorithm, 91–93, 107, 110, 180, 181, 183, 184, 192, 193, 208, 211–213, 215
 - model-improving iteration, 185, 187, 191, 193
 - model-improvement step, 191, 193, 209, 214
 - surrogate, 229
 - functional, 230
 - physical, 230

- Taylor, 5, 35, 36, 90, 92, 173–174, 178–181, 200, 230–231
using Kriging, 233, 240
using polynomial interpolation
 in the Frobenius minimum norm sense, 80–81
 in the minimum-norm sense, 74–76
 linear, 24
 of any degree, 37–38
using polynomial regression
 linear, 24–25
 of any degree, 58–59
using radial basis functions, 231–233, 240
- molecular geometry optimization, 5
multidisciplinary, 4
multiindices, 36
multiobjective optimization, 250
- natural basis (of monomials)
 definition, 36
- Nelder–Mead method, *see* methods
- Newton fundamental polynomials (NFPs)
 definition, 46
 use in DFO, 87, 208
- noise, 3, 5–11, 167–168, 171, 234
 consistency, 58, 68, 72
- noisy functions, *see* noise
noisy problems, *see* noise
- nonconvexity, 5, 6, 108, 149, 202, 246
nondifferentiability, 3, 136
nonmonotone, 136
nonsmoothness, 11, 58, 123–127, 132, 136, 247
- numerical comparisons, 10–11, *see also* data profiles
- optimality conditions
 first-order necessary for inequality constraints, 246
 first-order necessary for simple bounds, 243
- parallel, *see also* methods
 asynchronous, 136, 248
 difficulty of parallelization, 142
 ease of parallelization, 11, 136, 168
 environment, xii, 6, 116, 225
 parameter, *see also* simplex diameter; trust-region radius
 mesh size, 115, 118
 bounded, 129
 converging to zero, 135
 subsequence converging to zero, 121, 123
 step size, 115, 118, 164
 bounded, 129
 converging to zero, 135
 subsequence converging to zero, 121, 123
- pattern, *see also* integer lattices; mesh or grid
 generator, 127
- pivot polynomials, 101, 209–211, 216–218
- pivotal algorithms, 89, 99–109, 209–211
- poisedness, *see also* completing;
 condition number; improving;
 Lagrange polynomials;
 Newton fundamental polynomials
- Λ -poisedness
 for polynomial interpolation, 42
 for polynomial regression, 63
 for polynomial regression (strong), 65
 for underdetermined polynomial interpolation (minimum-norm sense), 78
 for underdetermined quadratic polynomial interpolation (in the minimum Frobenius norm sense), 82
 for polynomial interpolation
 linear, 24
 for polynomial regression
 linear, 25
- polling, *see also* poll step
 poll vectors or directions, 116

- polynomials, *see also* manifold; model;
natural basis (of monomials);
pivot
basis, 36
space of, 36
tail, 232
- positive, *see also* positive basis; positive
spanning set
dependence, 16
generators, 244–247
independence, 16
span, 15
- positive basis, *see also* positive spanning
set
definition, 16
examples of, 19–21, 30
maximal, 16
minimal, 16
uniform, 21
- positive spanning set
characterizations of, 17
cosine measure of, 22
definition, 16
implication of descent
characterization, 19
- radial basis functions (RBF), *see also*
global optimization; model;
uniqueness
conditionally positive definite, 232,
240
- regression, *see also* error bounds;
Lagrange polynomials; model;
poisedness; uniqueness
 ℓ_1 , 60
conditions, 25, 58
regularized, 70–71
- response surface methodology (RSM),
234–235
- sampling, *see also* completing;
improving; interpolation;
model; poisedness; regression;
starting; underdetermined
interpolation
D-optimality, 234
dense, 234
- design and analysis of computer
experiments (DACE), 234
- design of experiments, 234
Latin hypercubes, 234
orthogonal arrays, 234
- scaling
of matrix, 27, 28, 54, 79
system, 46
- serial
environment or machine, 5, 136,
168
- shifting and scaling
of sample set, 48–49, 68, 84, 90,
96–99, 103–104, 210, 217
- simplex
centered simplex gradient, 169
derivatives, 136, 169, 170
diameter of, 30
diameter of (converging to zero),
153
gradient (as gradient of
interpolation model), 33
gradient (as gradient of regression
model), 33
gradient (definition), 32
gradient (error bound), 33
Hessian (definition), 170
normalized volume of, 31
restarting, 160
set, 30
shape of, 30
starting, 150, 161
vertices of, 30
volume of, 30
- simulation, *see also* black box
PowerSpice, 4
- starting, *see also* starting simplex
feasible point, 242, 247
point, 120, 125, 126, 236, 239
sample set, 100–102, 107
- stencil
gradient, 33
radius, 33
set, 33
- step
Cauchy, 175–176

- criticality, 165, 183, 184, 192, 193, 210, 213, 218, 221, 224, 225
- eigenstep, 177–178
- poll step, *see also* mesh or grid
 - complete polling, 116, 131–132, 139, 157, 215
 - error bound (unsuccessful polling), 23, 122, 134
 - opportunistic polling, 116, 118, 119, 136
 - search step, *see also* mesh or grid
 - use of surrogates, 235–237
- stochastic optimization, 6
- stopping criterion, 8, 10, 120, 123, 138, 144, 152, 165, 182
- surrogate, *see also* model
 - rigorous optimization frameworks, 4, 235–240
- Taylor, *see* error bounds; model
- termination
 - early or premature, 5, 8, 242
- trust region, *see* methods
 - radius, 174
 - converging to zero, 188, 197
 - update, 178–179, 184, 193
 - subproblem, 175, 202–204, 220, 223, 238–239, 248–249
 - subproblem (wedge), 216, 221, 223
- underdetermined interpolation, *see also*
 - error bounds; Lagrange
 - polynomials; model;
 - poisedness; uniqueness
- conditions, 73
 - in the best subbasis sense, 74–76
- uniformly bounded
 - away from zero
 - cosine measure, 119, 122, 123
 - step size, 130
 - volume, 97
 - constant in norm equivalence, 49
 - independently of number of sample points, 70
 - model Hessian, 83, 186
 - number of improvement steps, 91, 96, 164, 165, 180, 181, 184, 193, 211, 212, 214, 215
 - number of iterations, 182, 187
 - number of unsuccessful iterations, 219, 220
 - poisedness constant, 179
- uniqueness, *see also* Lagrange
 - polynomials
 - interpolation polynomial, 38
 - lack of (in space-mapping models), 240
 - radial basis function model, 232
 - regression polynomial, 59
 - underdetermined minimum
 - Frobenius norm polynomial, 81, 87
- unisolvant, *see also* poisedness
 - set, 38
- wedge, *see also* methods; trust region
 - subproblem
 - constraint, 215–220