

The connectivity-profile of random increasing k -trees

Alexis Darrasse[†]
APR - LIP6
UPMC
75005 Paris France
alexis.darrasse@lip6.fr

Hsien-Kuei Hwang*
Institute of Statistical Science
Academia Sinica
Taipei 115 Taiwan
hkhwang@stat.sinica.edu.tw

Olivier Bodini[†]
APR - LIP6
UPMC
75005 Paris France
olivier.bodini@lip6.fr

Michèle Soria[†]
APR - LIP6
UPMC
75005 Paris France
michele.soria@lip6.fr

Abstract

Random increasing k -trees represent an interesting, useful class of strongly dependent graphs for which analytic-combinatorial tools can be successfully applied. We study in this paper a notion called connectivity-profile and derive asymptotic estimates for it; some interesting consequences will also be given.

1 Introduction

A k -tree is a graph reducible to a k -clique by successive removals of a vertex of degree k whose neighbors form a k -clique. This class of k -trees has been widely studied in combinatorics (for enumeration and characteristic properties [5, 29]), in graph algorithms (many NP-complete problems on graphs can be solved in polynomial time on k -trees [2]), and in many other fields where k -trees were naturally encountered (see [2]). By construction, vertices in such structures are remarkably close, reflecting a highly strong dependent graph structure, and they exhibit with no surprise the scale-free property [20], yet somewhat unexpectedly many properties of random k -trees can be dealt with by standard combinatorial, asymptotic and probabilistic tools, thus providing an important model of synergistic balance between mathematical tractability and the predictive power for practical-world complex networks.

While the term “ k -trees” is not very informative and may indeed be misleading to some extent, they stand out by their underlying tree structure, related to their recursive definition, which facilitates the analysis of the properties and the exploration of the structure. Indeed, for $k = 1$, k -

trees are just ordinary trees, and for $k \geq 2$ a bijection [11] between k -trees and a nontrivial simple family of trees can be explicitly defined, which we will describe briefly in this paper. To reduce ambiguity of terms, *except for “ k -trees”, all other occurrences of “trees” in this paper are ordinary trees without cycles.*

The process of generating a k -tree begins with a k -clique, which is itself a k -tree; then the k -tree grows by linking a new vertex to every vertex of an existing k -clique, and to these vertices only. The same process continues; see Figure 1 for an illustration. Such a simple process is reminiscent of several other models proposed in the literature such as k -DAGs [13], random circuits [3], preferential attachment [4, 7, 21], and many other models (see, for example, [6, 17, 25]). While the construction rule in each of these models is very similar, namely, linking a new vertex to k existing ones, the mechanism of choosing the existing k vertices differs from one case to another, resulting in very different topology and dynamics.

Restricting to the procedure of choosing a k -clique each time a new vertex is added, there are several variants of k -trees proposed in the literature depending on the modeling needs. So k -trees can be either labeled [5], unlabeled [22], increasing [32], planar [32], non-planar [5], or plane [26] (planar with a given embedding in the plane), etc.

For example, the family of random Apollonian networks, corresponding to planar 3-trees, has recently been employed as a model for complex networks [1, 32]. In these frameworks, since the exact topology of the real networks is difficult or even impossible to describe, one is often led to the study of models that present similarities to some observed properties such as the degree of a node and the distance between two nodes of the real structures.

For the purpose of this paper, we distinguish between

*Part of the work of this author was performed while visiting LIP6; he thanks the Laboratory for hospitality and support.

[†]This work was partially supported by ANR under the contract GAMMA, n°BLAN07-2-195422.




Figure 1: *The first few steps of generating a 3-tree and a 4-tree. Obviously, these graphs show the high connectivity of k -trees.*

two models of random labeled non-plane k -trees; by non-plane we mean that we consider these graphs as given by a set of edges (and not by its graphical representation):

- *random simply-generated k -trees*, which correspond to a uniform probability distribution on this class of labeled k -trees, and
- *random increasing k -trees*, where we consider the iterative generation process: at each time step, all existing k -cliques are equally likely to be selected and the new vertex is added with a label which is greater than the existing ones.

The two models are in good analogy to the simply-generated family of trees of Meir and Moon [24] marked specially by the functional equation $f(z) = z\Phi(f(z))$ for the underlying enumerating generating function, and the increasing family of trees of Bergeron et al. [10], characterized by the differential equation $f'(z) = \Phi(f(z))$. Very different stochastic behaviors have been observed for these families of trees. While similar in structure to these trees, the analytic problems on random k -trees we are dealing with here are however more involved because instead of a single scalar equation (either functional, algebraic, or differential), we now have a system of equations.

It is known that random trees in the family of increasing trees are often less skewed, less slanted in shape, a typical description being the logarithmic order for the distance of two randomly chosen nodes; this is in sharp contrast to the square-root order for random trees belonging to the simply-generated family; see for example [10, 14, 19, 23, 24]. Such a contrast has inspired and stimulated much recent research. Indeed, the majority of random trees in the literature of discrete probability, analysis of algorithms, and random combinatorial structures are either $\log n$ -trees or \sqrt{n} -trees, n being the tree size. While the class of \sqrt{n} -trees have been extensively investigated by probabilists and combinatorialists,

$\log n$ -trees are comparatively less addressed, partly because most of them were encountered not in probability or in combinatorics, but in the analysis of algorithms.

Table 1 presents a comparison of the two models: the classes \mathcal{T}_s and \mathcal{T} , corresponding respectively to simply-generated k -trees and increasing k -trees. The results concerning simple k -trees are given in [11, 12], and those concerning increasing k -trees are derived in this paper (except for the power law distribution [20]). We start with the specification, described in terms of operators of the symbolic method [18]. A structure of \mathcal{T}_s is a set of k structures of the same type, whose roots are attached to a new node: $\mathcal{T}_s = \text{Set}(\mathcal{Z} \times \mathcal{T}_s^k)$, while a structure of \mathcal{T} is an increasing structure, in the sense that the new nodes get labels that are smaller than those of the underlying structure (this constraint is reflected by the box-operator) $\mathcal{T} = \text{Set}(\mathcal{Z}^\square \times \mathcal{T}^k)$. The analytic difference immediately appears in the enumerative generating functions that translate the specifications: the simply-generated structures are defined by $T_s(z) = \exp(zT_s^k(z))$ and corresponding increasing structures satisfy the differential equation $T'(z) = T^k(z)$. These equations lead to a singular expansion of the square-root type in the simply-generated model, and a singularity in $(1 - kz)^{-1/k}$ in the increasing model. Similar analytic differences arise in the bivariate generating functions of shape parameters.

The expected distance between two randomly chosen vertices or the average path length is one of the most important shape parameters in modeling complex networks as it indicates roughly how efficient the information can be transmitted through the network. Following the same \sqrt{n} -vs- $\log n$ pattern, it is of order \sqrt{n} in the simply-generated model, but $\log n$ in the increasing model. Another equally important parameter is the degree distribution of a random vertex: its limiting distribution is a power law with exponential tails in the simply-generated model of the form $d^{-3/2}\rho_k^d$, in contrast to a power-law in the increasing model of the form $d^{-1-k/(k-1)}$, d denoting the degree [20]. As regards the de-

Model \ Properties	Simply-generated structures	Increasing structures
Combinatorial description	$\mathcal{T}_s = \text{Set}(\mathcal{Z} \times \mathcal{T}_s^k)$	$\mathcal{T} = \text{Set}(\mathcal{Z}^\square \times \mathcal{T}^k)$
Generating function	$T_s(z) = \exp(zT_s^k(z))$	$T'(z) = T^k(z)$
Expansion near singularity	$T_s(z) = \tau - h\sqrt{1 - z/\rho} + \dots$	$T(z) = (1 - kz)^{-1/k}$
Mean distance of nodes	$O(\sqrt{n})$	$O(\log n)$
Degree distribution	Power law with exp. tails	Power law [20]
Root-degree distribution	Power law with exp. tails	Stable law (Theorem 4.1)
Expected Profile	Rayleigh limit law	Gaussian limit law (3.8)

Table 1: The contrast of some properties between random simply-generated k -trees and random increasing k -trees. Here \mathcal{Z} denotes a node and \mathcal{Z}^\square means a node with the smallest label.

gree of the root-vertex¹, its asymptotic distribution remains the same as that of any vertex in the simply-generated model, but in the increasing model, the root-degree distribution is different, with an asymptotic stable law (which is Rayleigh in the case $k = 2$); see Theorem 4.1.

Our main concern in this paper is the connectivity-profile. Recall that the profile of an usual tree is the sequence of numbers, each enumerating the total number of nodes with the same distance to the root. For example, the tree $\leftarrow\leftarrow\leftarrow\leftarrow$ has the profile $\{1, 2, 2, 1, 3\}$. Profiles represent one of the richest shape measures and they convey much information regarding particularly the silhouette. On random trees, they have been extensively studied recently; see [8, 15, 16, 19, 21, 23, 27]. Since k -trees have many cycles for $k \geq 2$, we call the profile of the shortest-path tree (see next section) *the connectivity-profile* as it measures to some extent the connectivity of the graph on the one hand and is well connected to the notion of profiles of trees on the other hand. Indeed this connectivity-profile corresponds to the profile of the “shortest-path tree” of a k -tree, as defined by Proskurowski [28], which is nothing more than the result of a Breadth First Search (BFS) on the graph. Moreover, in the domain of complex networks, this kind of BFS trees is an important object; for example, it describes the results of the `traceroute` measuring tool [30, 31] in the study of the topology of the Internet.

We will derive precise asymptotic approximations to the expected connectivity-profile of random increasing k -trees, the major tools used being based on the resolution of a system of differential equations of Cauchy-Euler type (see [9]). In particular, the expected number of nodes at distance d from the root-vertex follows asymptotically a Gaussian distribution, in contrast to the Rayleigh limit

distribution in the case of simply-generated k -trees. Also the limit distribution of the number of nodes with distance d to the root-vertex will be derived when d is bounded. Note that when $d = 1$, the number of nodes at distance 1 to the root-vertex is nothing but the degree of the root-vertex.

This paper is organized as follows. We first present the definition and combinatorial specification of random increasing k -trees in Section 2, together with the enumerative generating functions, on which our analytic tools will be based. We then present two asymptotic approximations to the expected connectivity-profile in Section 3, one for $d = o(\log n)$ and the other for $d \rightarrow \infty$ and $d = O(\log n)$. Interesting consequences of our results will also be given. The limit distribution of the connectivity-profile in the range when $d = O(1)$ is then given in Section 4.

2 Random increasing k -trees and generating functions

Since k -trees are graphs full of cycles and cliques, the key step in our analytic-combinatorial approach is to introduce a bijection between k -trees and a suitably defined class of trees for which generating functions can be derived. This approach was successfully applied to simply-generated family of k -trees in [11], which leads to a system of algebraic equations. The bijection argument used there can be adapted *mutatis mutandis* here for increasing k -trees, which then yields a system of differential equations through the bijection with a class of increasing trees [10].

Increasing k -trees and the bijection. Recall that a k -clique is a set of k mutually adjacent vertices.

DEFINITION 2.1. *An increasing k -tree is defined recursively as follows. A k -clique in which each vertex gets a distinct label from $\{1, \dots, k\}$ is an increasing k -tree of k vertices. An increasing k -tree with $n > k$ vertices is constructed from an increasing k -tree with $n - 1$ vertices by adding a vertex labeled n and by connecting it by an edge to each of the k vertices in an existing k -clique.*

¹We call *root-clique* the clique composed of the k vertices $(1, \dots, k)$. The increasing nature of the k -trees guarantees that these vertices always form a clique. We call *root-vertex* the vertex with label 1.




Figure 2: A 2-tree (left) and its corresponding increasing tree representation (right).

By *random increasing k -trees*, we assume that all existing k -cliques are equally likely each time a new vertex is being added. Given the fact that each time a vertex is added, k new cliques appear in the random increasing k -tree, it is immediate that the number T_n of increasing k -trees of $n + k$ nodes is given by $T_n = \prod_{0 \leq i < n} (ik + 1)$.

Note that if we allow any permutation on all labels, we obtain the class of simply-generated k -trees where monotonicity of labels along paths fails in general.

Combinatorially, simply-generated k -trees are in bijection [11] with the family of trees specified by $\mathcal{K}_s = \mathcal{Z}^k \times \mathcal{T}_s$, where $\mathcal{T}_s = \text{Set}(\mathcal{Z} \times \mathcal{T}_s^k)$. Given a rooted k -tree G of n vertices, we can transform G into a tree T , with the root node labeled $\{1, \dots, k\}$, by the following procedure. First, associate a white node to each k -clique of G and a black node to each $(k + 1)$ -clique of G . Then add a link between each black node and all white nodes associated to the k -cliques it contains. Each black node is labeled with the only vertex not appearing in one of the black nodes above it or in the root. The last step in order to complete the bijection is to order the k vertices of the root and propagate this order to the k sons of each black node. This constructs a tree from a k -tree (see Figure 2); conversely, we can obtain the k -tree through a simple traversal of the tree.

Such a bijection translates directly to increasing k -trees by restricting the class of corresponding trees to those respecting a monotonicity constraint on the labels, namely, on any path from the root to a leaf the labels are in increasing order. This yields the combinatorial specification of the class of increasing trees $\mathcal{T} = \text{Set}(\mathcal{Z}^\square \times \mathcal{T}^k)$. An increasing k -tree is just a tree in \mathcal{T} together with the sequence $\{1, \dots, k\}$ corresponding to the labels of the root-clique. A tree in $\mathcal{K} = \mathcal{Z}^\square \times \dots \times \mathcal{Z}^\square \times \mathcal{T}$ is thus completely determined by its \mathcal{T} component, giving $\mathcal{K}_{n+k} \equiv \mathcal{T}_n$. For example figure 2 shows a 2-tree with 19 vertices and its tree representation with 17 black nodes. In the rest of this paper we will thus focus on class \mathcal{T} .

Generating functions. Following the bijection, we see that the complicated dependence structure of k -trees is now completely described by the class of increasing trees specified by $\mathcal{T} = \text{Set}(\mathcal{Z}^\square \times \mathcal{T}^k)$. Let $T(z) := \sum_{n \geq 0} T_n z^n / n!$ denote the exponential generating function of the number T_n of increasing k -trees of $n + k$ vertices. Then, given that the box construction $\mathcal{Z}^\square \times \mathcal{A}$ translates into the primitive $\int_0^z A(x) dx$, the specification translates into the equation

$$T(z) = \exp \left(\int_0^z T^k(x) dx \right),$$

or, equivalently, $T'(z) = T^{k+1}(z)$ with $T(0) = 1$, which is solved to be

$$T(z) = (1 - kz)^{-1/k},$$

we then check that $T_n = \prod_{0 \leq i < n} (ik + 1)$.

Marking by u the neighbors of the root-vertex in \mathcal{T} , we notice that one amongst k of the subtrees is special: it does not contain any neighbor of the root-vertex and the $k - 1$ others have the same structure as the whole tree. We thus obtain

$$T(z, u) = \exp \left(u \int_0^z T(x) T^{k-1}(x, u) dx \right),$$

where the coefficients $n! [u^\ell z^n] T(z, u)$ denote the number of increasing k -trees of size $n + k$ with root degree equal to $k + \ell - 1$ (counting the $k - 1$ neighbors of the root-vertex coming from the root-clique). Taking derivative with respect to z on both sides and then solving the equation, we get the closed-form expression

$$(2.1) \quad T(z, u) = \left(1 - u(1 - (1 - kz)^{-1/k}) \right)^{-1/(k-1)}.$$

Since k -trees can be transformed into ordinary increasing trees, the profiles of the shortest-path trees can be naturally defined, although they do not correspond to simple parameters on k -trees. Also the results provide more insight on the structure of random k -trees. Roughly, we expect that

all vertices on k -trees are close, one at most of logarithmic order away from the other. The fine results we derive provide in particular an upper bound for that.

Let $X_{n;d,j}$ denote the number of nodes at distance d from j vertices of the root-clique in a random k -tree of $n+k$ vertices. Let $T_{d,j}(z, u) = \sum_{n \geq 0} T_n \mathbb{E}(u^{X_{n;d,j}}) z^n / n!$ denote the corresponding bivariate generating function.

THEOREM 2.1. *The generating functions $T_{d,j}$'s satisfy the differential equations*

$$(2.2) \quad \frac{\partial}{\partial z} T_{d,j}(z, u) = u^{\delta_{d,1}} T_{d,j-1}^j(z, u) T_{d,j}^{k-j+1}(z, u),$$

with the initial conditions $T_{d,j}(0, u) = 1$ for $1 \leq j \leq k$, where $\delta_{a,b}$ denotes the Kronecker function, $T_{0,k}(z, u) = T(z)$ and $T_{d,0}(z, u) = T_{d-1,k}(z, u)$.

Proof. The theorem follows from

$$T_{d,j}(z, u) = \exp\left(u^{\delta_{d,1}} \int_0^z T_{d,j-1}^j(x, u) T_{d,j}^{k-j}(x, u) dx\right),$$

with $T_{d,j}(z, 1) = T(z)$.

For operational convenience, we normalize all z by z/k and write $\tilde{T}(z) := T(z/k) = (1-z)^{-1/k}$. Similarly, we define $\tilde{T}_{d,j}(z, u) := T_{d,j}(z/k, u)$ and have, by (2.2),

$$(2.3) \quad \frac{\partial}{\partial z} \tilde{T}_{d,j}(z, u) = \frac{u^{\delta_{d,1}}}{k} \tilde{T}_{d,j-1}^j(z, u) \tilde{T}_{d,j}^{k-j+1}(z, u),$$

with $\tilde{T}_{d,j}(1, z) = \tilde{T}(z)$, $\tilde{T}_{0,k}(z, u) = \tilde{T}(z)$ and $\tilde{T}_{d,0}(z, u) = \tilde{T}_{d-1,k}(z, u)$.

3 Expected connectivity-profile

We consider the expected connectivity-profile $\mathbb{E}(X_{n;d,j})$ in this section. Observe first that

$$\mathbb{E}(X_{n;d,j}) = \frac{k^n [z^n] \tilde{M}_{d,j}(z)}{T_n},$$

where $\tilde{M}_{d,j}(z) := \partial \tilde{T}_{d,j}(z, u) / (\partial u)|_{u=1}$. It follows from (2.3) that

$$(3.4) \quad \begin{aligned} \tilde{M}'_{d,j}(z) &= \frac{(k-j+1)\tilde{M}_{d,j}(z) + j\tilde{M}_{d,j-1}(z)}{k(1-z)} \\ &\quad + \frac{\delta_{d,1}\tilde{T}(z)}{k(1-z)}. \end{aligned}$$

This is a standard differential equation of Cauchy-Euler type whose solution is given by (see [9])

$$\begin{aligned} \tilde{M}_{d,j}(z) &= \frac{(1-z)^{-(k-j+1)/k}}{k} \int_0^z (1-x)^{-(j-1)/k} \\ &\quad \times \left(j\tilde{M}_{d,j-1}(x) + \delta_{d,1}\tilde{T}(x) \right) dx, \end{aligned}$$

since $\tilde{M}_{d,j}(0) = 0$. Then, starting from $\tilde{M}_{0,k} = 0$, we get

$$\tilde{M}_{1,1}(z) = \frac{1}{k-1} \left(\frac{1}{1-z} - \frac{1}{(1-z)^{1/k}} \right).$$

And by induction, we get

$$\tilde{M}_{d,j}(z) \sim \frac{j}{(k-1)(d-1)!} \cdot \frac{1}{1-z} \log^{d-1} \frac{1}{1-z},$$

for $1 \leq j \leq k$, $d \geq 1$ and $z \sim 1$. One can then prove, by singularity analysis, that

$$\mathbb{E}(X_{n;d,j}) \sim \Gamma(1/k) \frac{j}{k-1} \cdot \frac{(\log n)^{d-1}}{(d-1)!} n^{1-1/k},$$

for large n and fixed d, k and $1 \leq j \leq k$. We can indeed prove that the same asymptotic estimate holds in a larger range.

THEOREM 3.1. *The expected connectivity-profile $\mathbb{E}(X_{n;d,j})$ satisfies for $1 \leq d = o(\log n)$*

$$(3.5) \quad \mathbb{E}(X_{n;d,j}) \sim \Gamma(1/k) \frac{j}{k-1} \cdot \frac{(\log n)^{d-1}}{(d-1)!} n^{1-1/k},$$

uniformly in d , and for $d \rightarrow \infty$, $d = O(\log n)$,

$$(3.6) \quad \mathbb{E}(X_{n;d,j}) \sim \frac{\Gamma(1/k) h_{j,1}(\rho) \rho^{-d} n^{\lambda_1(\rho)-1/k}}{\Gamma(\lambda_1(\rho)) \sqrt{2\pi(\rho \lambda_1'(\rho) + \rho^2 \lambda_1''(\rho)) \log n}}$$

where $\rho = \rho_{n,d} > 0$ solves the equation $\rho \lambda_1'(\rho) = d / \log n$, $\lambda_1(w)$ being the largest zero (in real part) of the equation $\prod_{1 \leq \ell \leq k} (\theta - \ell/k) - k!w/k^k = 0$ and satisfies $\lambda_1(1) = (k+1)/k$, and $h_{j,1}$ is defined below.

An explicit expression for the $h_{j,1}$'s is given as follows. Let $\lambda_1(w), \dots, \lambda_k(w)$ denote the zeros of the equation $\prod_{1 \leq \ell \leq k} (\theta - \ell/k) - k!w/k^k = 0$. Then for $1 \leq j \leq k$

$$(3.7) \quad \begin{aligned} h_{j,1}(w) &= \frac{j!w(w-1)}{(k\lambda_1(w)-1) \prod_{k-j+1 \leq s \leq k+1} (k\lambda_1(w)-s)} \\ &\quad \times \frac{1}{\sum_{1 \leq s \leq k} \frac{1}{k\lambda_1(w)-s}}. \end{aligned}$$

The theorem cannot be proved by the above inductive argument and our method of proof consists of the following steps. First, the bivariate generating functions $\mathcal{M}_j(z, w) := \sum_{d \geq 1} \tilde{M}_{d,j}(z) w^d$ satisfy the linear system

$$\left((1-z) \frac{d}{dz} - \frac{k-j+1}{k} \right) \mathcal{M}_j = \frac{j}{k} \mathcal{M}_{j-1} + \frac{w\tilde{T}}{k},$$

for $1 \leq j \leq k$. Second, this system is solved and has the solutions

$$\mathcal{M}_j(z, w) = \sum_{1 \leq j \leq k} h_{j,m}(w)(1-z)^{-\lambda_m(w)} - \frac{w - (w-1)\delta_{k,j}}{k} \tilde{T}(z),$$

where the $h_{j,m}$ have the same expression as $h_{j,1}$ but with all $\lambda_1(w)$ in (3.7) replaced by $\lambda_m(w)$. While the form of the solution is well anticipated, the hard part is the calculations of the coefficient-functions $h_{j,m}$. Third, by singularity analysis and a delicate study of the zeros, we then conclude, by saddle-point method, the estimates given in the theorem.

COROLLARY 3.1. *The expected degree of the root-vertex $\mathbb{E}(X_{n,1,j})$ satisfies*

$$\mathbb{E}(X_{n,1,j}) \sim \Gamma(1/k) \frac{j}{k-1} n^{1-1/k} \quad (1 \leq j \leq k).$$

This estimate also follows easily from (2.1).

Let $H_k := \sum_{1 \leq \ell \leq k} 1/\ell$ denote the harmonic numbers and $H_k^{(2)} := \sum_{1 \leq \ell \leq k} 1/\ell^2$.

COROLLARY 3.2. *The expected number of nodes at distance $d = \lfloor \frac{1}{kH_k} \log n + x\sigma\sqrt{\log n} \rfloor$ from the root-vertex, where $\sigma = \sqrt{H_k^{(2)}/(kH_k^3)}$, satisfies, uniformly for $x = o((\log n)^{1/6})$,*

$$(3.8) \quad \mathbb{E}(X_{n,d,j}) \sim \frac{ne^{-x^2/2}}{\sqrt{2\pi\sigma^2 \log n}}.$$

This Gaussian approximation justifies the last item corresponding to increasing trees in Table 1.

Note that $\lambda_1(1) = (k+1)/k$ and $\alpha = d/\log n \sim 1/(kH_k)$. In this case, $\rho = 1$ and

$$\rho\lambda_1'(\rho) = \frac{1}{\sum_{1 \leq \ell \leq k} \frac{1}{\lambda_1(\rho) - \frac{\ell}{k}}},$$

which implies that $\lambda_1(\rho) - 1/k - \alpha \log \rho \sim 1$.

COROLLARY 3.3. *Let $\mathcal{H}_{n;d,j} := \max_d X_{n;d,j}$ denote the height of a random increasing k -tree of $n+k$ vertices. Then*

$$\mathbb{E}(\mathcal{H}_n) \leq \alpha_+ \log n - \frac{\alpha_+}{2(\lambda_1(\alpha_+) - \frac{1}{k})} \log \log n + O(1),$$

where $\alpha_+ > 0$ is the solution of the system of equations

$$\begin{cases} \frac{1}{\alpha_+} = \sum_{1 \leq \ell \leq k} \frac{1}{v - \frac{\ell}{k}}, \\ v - \frac{1}{k} - \alpha_+ \sum_{1 \leq \ell \leq k} \log \left(\frac{k}{\ell} v - 1 \right) = 0. \end{cases}$$

Table 2 gives the numerical values of α_+ for small values of k . For large k , one can show that $\alpha_+ \sim 1/(k \log 2)$ and $\lambda_1(\alpha_+) \sim 2$.

Corollary 3.3 justifies that the mean distance of random k -trees are of logarithmic order in size, as stated in Table 1.

COROLLARY 3.4. *The width $\mathcal{W}_{n;d,j} := \max_d X_{n;d,j}$ is bounded from below by*

$$\mathbb{E}(\mathcal{W}_n) = \mathbb{E}(\max_d X_{n,d}) \geq \max_d \mathbb{E}(X_{n,d}) \sim c \frac{n}{\sqrt{\log n}}.$$

We may conclude briefly from all these results that *in the shortest-path increasing trees of random increasing k -trees, almost all nodes are located in the levels with $d = \frac{1}{kH_k} \log n + O(\sqrt{\log n})$, each with $n/\sqrt{\log n}$ nodes.*

4 Limiting distributions

With the availability of the bivariate generating functions (2.2), we can proceed further and derive the limit distribution of $X_{n;d,j}$ in the range where $d = O(1)$. The case when $d \rightarrow \infty$ is much more involved; we content ourselves in this extended abstract with the statement of the result for bounded d .

THEOREM 4.1. *The random variables $X_{n;d,j}$, when normalized by their mean orders, converge in distribution to*

$$(4.9) \quad \frac{X_{n;d,j}}{n^{1-1/k}(\log n)^{d-1}/(d-1)!} \xrightarrow{d} \Xi_{d,j},$$

where

$$\begin{aligned} \mathbb{E}(e^{\Xi_{d,j}u}) &= \Gamma\left(\frac{1}{k}\right) \sum_{m \geq 0} \frac{c_{d,j,m}}{m! \Gamma(m(1-1/k) + 1/k)} u^m \\ &= \frac{\Gamma\left(\frac{1}{k}\right)}{2\pi i} \int_{-\infty}^{(0+)} e^{\tau} \tau^{-1/k} C_{d,j} \left(\tau^{-1+1/k} u \right) d\tau, \end{aligned}$$

and $C_{d,j}(u) := 1 + \sum_{m \geq 1} c_{d,j,m} u^m / m!$ satisfies the system of differential equations

$$(4.10) \quad (k-1)uC_{d,j}'(u) + C_{d,j}(u) = C_{d,j}(u)^{k+1-j} C_{d,j-1}(u)^j,$$

for $1 \leq j \leq k$, with $C_{d,0} = C_{d-1,k}$. Here the symbol $\int_{-\infty}^{(0+)}$ denotes any Hankel contour starting from $-\infty$ on the real axis, encircling the origin once counter-clockwise, and returning to $-\infty$.

We indeed prove the convergence of all moments, which is stronger than weak convergence; also the limit law is uniquely determined by its moment sequence.

So far only in special cases do we have explicit solution for $C_{1,j}$: $C_{1,1}(u) = (1+u)^{-1/(k-1)}$ and

$$C_{1,2}(u) = \begin{cases} \frac{e^{1/(1+u)}}{1+u}, & \text{if } k = 2; \\ \frac{1}{1 + u^{1/2} \arctan(u^{1/2})}, & \text{if } k = 3. \end{cases}$$

k	2	3	4	5	6
α_+	1.085480	0.656285	0.465190	0.358501	0.290847
k	7	8	9	10	20
α_+	0.244288	0.210365	0.184587	0.164356	0.077875

Table 2: Approximate numerical values of α_+ .

Note that the result (4.9) when $d = 0$ can also be derived directly by the explicit expression (2.1). In particular, when $k = 2$, the limit law is Rayleigh.

References

- [1] Jr. Andrade, Hans J. Herrmann, Roberto F. S. Andrade, and Luciano R. da Silva. Apollonian networks: Simultaneously Scale-Free, small world, euclidean, space filling, and with matching graphs. *Physical Review Letters*, 94(1):018702–4, 2005.
- [2] Stefan Arnborg, Derek G. Corneil, and Andrzej Proskurowski. Complexity of finding embeddings in a k -tree. *SIAM Journal on Algebraic and Discrete Methods*, 8(2):277–284, April 1987.
- [3] Sunil Arya, Mordecai J. Golin, and Kurt Mehlhorn. On the expected depth of random circuits. *Combinatorics, Probability and Computing*, 8(03):209–228, 1999.
- [4] Albert-László Barabási and Réka Albert. Emergence of scaling in random networks. *Science*, 286(5439):509–512, October 1999.
- [5] Lowell W. Beineke and Raymond E. Pippert. The number of labeled k -dimensional trees. *Journal of Combinatorial Theory*, 6(2):200–205, March 1969.
- [6] S. Boccaletti, V. Latora, Y. Moreno, M. Chavez, and D.-U. Hwang. Complex networks: Structure and dynamics. *Physics Reports*, 424(4-5):175–308, February 2006.
- [7] Béla Bollobás, Oliver Riordan, Joel Spencer, and Gábor Tusnády. The degree sequence of a scale-free random graph process. *Random Structures and Algorithms*, 18(3):279–290, 2001.
- [8] Brigitte Chauvin, Thierry Klein, Jean-François Marckert, and Alain Rouault. Martingales and profile of binary search trees. *Electronic Journal of Probability*, 10:420–435, June 2005.
- [9] Hua-Huai Chern, Hsien-Kuei Hwang, and Tsung-Hsi Tsai. An asymptotic theory for Cauchy-Euler differential equations with applications to the analysis of algorithms. *Journal of Algorithms*, 44(1):177–225, July 2002.
- [10] François Bergeron, Philippe Flajolet, and Bruno Salvy. Varieties of increasing trees. In *CAAP '92*, pages 24–48. 1992.
- [11] Alexis Darrasse and Michèle Soria. Limiting distribution for distances in k -trees. In *Proceedings of the 20th International Workshop on Combinatorial Algorithms (IWOCA 2009)*, Lecture Notes in Computer Science. Springer-Verlag, June 2009.
- [12] Alexis Darrasse, Michèle Soria, and Olivier Bodini. A unifying structural approach to the analysis of parameters in k -trees. *In preparation*.
- [13] Luc Devroye and Svante Janson. Long and short paths in uniform random recursive dags. *arXiv:0906.0152v1*, May 2009.
- [14] Michael Drmota. *Random Trees*. Springer Vienna, April 2009.
- [15] Michael Drmota and Bernhard Gittenberger. On the profile of random trees. *Random Structures and Algorithms*, 10(4):421–451, 1997.
- [16] Michael Drmota, Svante Janson, and Ralph Neininger. A functional limit theorem for the profile of search trees. *The Annals of Applied Probability*, 18(1):288–333, 2008.
- [17] Rick Durrett. *Random Graph Dynamics*. Number 20 in Cambridge Series in Statistical and Probabilistic Mathematics. Cambridge University Press, 2006.
- [18] Philippe Flajolet and Robert Sedgewick. *Analytic Combinatorics*. Cambridge University Press, 2009.
- [19] Michael Fuchs, Hsien-Kuei Hwang, and Ralph Neininger. Profiles of random trees: Limit theorems for random recursive trees and binary search trees. *Algorithmica*, 46(3):367–407, November 2006.
- [20] Yong Gao. The degree distribution of random k -trees. *Theoretical Computer Science*, 410(8-10):688–695, March 2009.
- [21] Hsien-Kuei Hwang. Profiles of random trees: Plane-oriented recursive trees. *Random Structures and Algorithms*, 30(3):380–413, 2007.
- [22] Gilbert Labelle, Cédric Lamathe, and Pierre Leroux. Labeled and unlabelled enumeration of k -gonal 2-trees. *Journal of Combinatorial Theory, Series A*, 106(2):193–219, May 2004.
- [23] Jean-François Marckert and Marie Albenque. Some families of increasing planar maps. *Electronic Journal of Probability*, 13:1624–1671, September 2008.
- [24] Amram Meir and John W. Moon. On the altitude of nodes in random trees. *Canadian Journal of Mathematics*, 30:997–1015, 1978.
- [25] Mark E. J. Newman. The structure and function of complex networks. *SIAM Review*, 45(2):167–256, 2003.
- [26] Edgar M. Palmer and Ronald C. Read. On the number of plane 2-Trees. *J. London Math. Soc.*, s2-6(4):583–592, 1973.
- [27] Gahyun Park, Hsien-Kuei Hwang, Pierre Nicodème, and Wojciech Szpankowski. Profiles of tries. *SIAM Journal on Computing*, 38(5):1821, 2009.
- [28] Andrzej Proskurowski. K -trees: representation and distances. In *Congressus Numerantium*, volume 29, pages 785–794. Utilitas Mathematica, 1980.
- [29] Donald J. Rose. On simple characterizations of k -trees. *Discrete Mathematics*, 7(3-4):317–322, 1974.
- [30] W. Richard Stevens. Chapter 8, traceroute program. In

TCP/IP Illustrated, Volume 1: The Protocols. Addison-Wesley Professional, 1994.

- [31] Fabien Viger, Brice Augustin, Xavier Cuvellier, Clémence Magnien, Matthieu Latapy, Timur Friedman, and Renata Teixeira. Detection, understanding, and prevention of traceroute measurement artifacts. *Computer Networks*, 52(5):998–1018, April 2008.
- [32] Zhongzhi Zhang, Lili Rong, and Francesc Comellas. High-dimensional random apollonian networks. *Physica A: Statistical Mechanics and its Applications*, 364:610–618, May 2006.